

RESEARCH OUTPUTS / RÉSULTATS DE RECHERCHE

Draft Genome Sequences of Three Capnocytophaga cynodegmi Strains Isolated from the Oral Cavity of Healthy Dogs

Manfredi, Pablo; Renzi, Francesco; Cornelis, Guy

Published in:
Genome Announcements

DOI:
[10.1128/genomeA.00200-15](https://doi.org/10.1128/genomeA.00200-15)

Publication date:
2015

Document Version
Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for pulished version (HARVARD):
Manfredi, P, Renzi, F & Cornelis, G 2015, 'Draft Genome Sequences of Three Capnocytophaga cynodegmi Strains Isolated from the Oral Cavity of Healthy Dogs', *Genome Announcements*.
<https://doi.org/10.1128/genomeA.00200-15>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Draft Genome Sequences of Three *Capnocytophaga cynodegmi* Strains Isolated from the Oral Cavity of Healthy Dogs

Pablo Manfredi,^a Francesco Renzi,^{a,b} Guy R. Cornelis^{a,b}

Biozentrum der Universität Basel, Basel, Switzerland^a; University of Namur, Namur, Belgium^b

Here, we present the draft genome sequences of three strains of *Capnocytophaga cynodegmi*. In contrast to the very close relationship among them, *C. cynodegmi* and *Capnocytophaga canimorsus* differ dramatically in terms of virulence in humans. Comparative genomics provided some understanding on how *Capnocytophaga* species may switch from being dog commensals to human pathogens.

Received 12 February 2015 Accepted 28 April 2015 Published 28 May 2015

Citation Manfredi P, Renzi F, Cornelis GR. 2015. Draft genome sequences of three *Capnocytophaga cynodegmi* strains isolated from the oral cavity of healthy dogs. *Genome Announc* 3(3):e00200-15. doi:10.1128/genomeA.00200-15.

Copyright © 2015 Manfredi et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Guy R. Cornelis, guy.cornelis@unamur.be.

Capnocytophaga cynodegmi (formerly CDC dysgonic fermenter-2-like) is a common oral commensal of dogs and cats, with prevalence rates as high as 86% and 84%, respectively (1). It belongs to the *Bacteroidetes* phylum, where it is very closely related to *Capnocytophaga canimorsus*, another commensal of the dog mouth (2, 3). Together, the two species display a significant number of features that differentiate them from other members of the *Capnocytophaga* genus (4). Beside subtle differences in their 16S rRNA gene sequences (4), *C. cynodegmi* strains can be differentiated from *C. canimorsus* strains by the light-yellow color displayed by colonies grown on sheep blood agar and by the capacity to ferment sucrose, raffinose, inulin, and melibiose (4). Contrary to *C. canimorsus*, *C. cynodegmi* is not a severe and sepsis-causing human pathogen, although some cases of human infections have been reported to consist mostly of wound or corneal infections (4, 5–8).

Three strains of *C. cynodegmi*, Ccy74, Ccyn2B, and Ccyn_ATCC 49044, were isolated from canine oral swabs and identified by 16S rRNA sequencing (4, 9). The strains were selected as dispersed representatives of the species *C. cynodegmi*, according to 16S rRNA phylogenetics and limited phenotyping (9). Genomic DNA was extracted using the Genomic-tip 500/G DNA extraction kit (catalog no. 10262; Qiagen), according to the manufacturer's instructions, followed by an additional phenol-chloroform purification step. Sequencing was performed at LGC Genomics, Berlin, Germany, on one Illumina HiSeq 2000 channel and generated approximately 10.8 ± 1.0 million 100-bp single reads per strain. *De novo* assembly was performed with Velvet, with optimized parameters (10). On average, draft assemblies accounted for 2.69 ± 0.01 Mb for 90 (Ccy74), 67 (Ccyn2B), and 111 (Ccyn_ATCC 49044) contigs. Genome metrics and automated annotation were conducted at the LABGeM, France Génomique (11). The G+C content ($34.40\% \pm 0.01\%$) is lower than that of the closely related *C. canimorsus* ($36.16\% \pm 0.08\%$). Each genome contains a fairly similar number of coding sequences ($2,484 \pm 18$). Ccyn2B exhibits significantly more strain-specific coding sequences (CDSs) (350) than those in Ccy74 (75) and Ccyn_ATCC 49044 (72).

The *C. cynodegmi* core genome is composed of 1,910 families of orthologs, of which 341 are specific to *C. cynodegmi* compared to seven genomes of *C. canimorsus* (2, 3, 12). While 253 clusters of orthologs were of unknown function, genes involved in aromatic amino acid synthesis (the complete L-tryptophan synthesis pathway from chorismate, 5 genes), glycan chain foraging (a complete polysaccharide utilization locus, 8 genes) (13), oxidative respiration and oxidative stress resistance (5 genes), and lipopolysaccharide (LPS) and polysaccharide biosynthesis (5 genes) formed the major functional clusters of the species-exclusive core genome. With respect to iron acquisition, a homolog to the heme-binding HmuY protein (14) was found in each of the three *C. cynodegmi* genomes, in addition to a locus encoding the iron capture system of *Bacteroidetes* (15).

Nucleotide sequence accession numbers. These whole-genome shotgun projects have been deposited in ENA under the accession numbers [CDOD00000000](https://ena.ebi.ac.uk/ena/record/CDOD00000000) (Ccyn2B), [CDOG00000000](https://ena.ebi.ac.uk/ena/record/CDOG00000000) (Ccy74), and [CDOF00000000](https://ena.ebi.ac.uk/ena/record/CDOF00000000) (Ccyn_ATCC 49044). The versions described in this paper are the initial versions.

ACKNOWLEDGMENTS

This work was supported by grants 3100A0-128659 from the Swiss National Science Foundation and ERC 2011-ADG 20110310 from the European Research Council.

F.R. is a postdoctoral research fellow (chargé de recherche) of the Belgian Fonds National de la Recherche Scientifique (FNRS).

REFERENCES

1. Suzuki M, Kimura M, Imaoka K, Yamada A. 2010. Prevalence of *Capnocytophaga canimorsus* and *Capnocytophaga cynodegmi* in dogs and cats determined by using a newly established species-specific PCR. *Vet Microbiol* 144:172–176. <http://dx.doi.org/10.1016/j.vetmic.2010.01.001>.
2. Manfredi P, Renzi F, Cornelis GR. 2015. Draft genome sequences of three *Capnocytophaga canimorsus* strains isolated from healthy canine oral cavities. *Genome Announc* 3(3):e00199–15. <http://dx.doi.org/10.1128/genomeA.00199-15>.
3. Manfredi P, Renzi F, Cornelis GR. 2015. Draft genome sequences of three *Capnocytophaga canimorsus* strains isolated from septic patients. *Genome Announc* 3(3):e00193–15. <http://dx.doi.org/10.1128/genomeA.00193-15>.

4. Brenner DJ, Hollis DG, Fanning GR, Weaver RE. 1989. *Capnocytophaga canimorsus* sp. nov. (formerly CDC group DF-2), a cause of septicemia following dog bite, and *C. cynodegmi* sp. nov., a cause of localized wound infection following dog bite. J Clin Microbiol 27:231–235.
5. Le Moal G, Landron C, Grollier G, Robert R, Burucoa C. 2003. Meningitis due to *Capnocytophaga canimorsus* after receipt of a dog bite: case report and review of the literature. Clin Infect Dis 36:e42–e46. <http://dx.doi.org/10.1086/345477>.
6. Lion C, Escande F, Burdin JC. 1996. *Capnocytophaga canimorsus* infections in human: review of the literature and cases report. Eur J Epidemiol 12:521–533. <http://dx.doi.org/10.1007/BF00144007>.
7. Sarma PS, Mohanty S. 2001. *Capnocytophaga cynodegmi* cellulitis, bacteremia, and pneumonitis in a diabetic man. J Clin Microbiol 39: 2028–2029. <http://dx.doi.org/10.1128/JCM.39.5.2028-2029.2001>.
8. Forman MA, Johnson LR, Jang S, Foley JE. 2005. Lower respiratory tract infection due to *Capnocytophaga cynodegmi* in a cat with pulmonary carcinoma. J Feline Med Surg 7:227–231. <http://dx.doi.org/10.1016/j.j-fms.2004.11.002>.
9. Mally M, Paroz C, Shin H, Meyer S, Soussoula LV, Schmiediger U, Saillen-Paroz C, Cornelis GR. 2009. Prevalence of *Capnocytophaga canimorsus* in dogs and occurrence of potential virulence factors. Microbes Infect 11:509–514. <http://dx.doi.org/10.1016/j.micinf.2009.02.005>.
10. Zerbino DR, Birney E. 2008. Velvet: algorithms for *de novo* short read assembly using de Bruijn graphs. Genome Res 18:821–829. <http://dx.doi.org/10.1101/gr.074492.107>.
11. Vallenet D, Belda E, Calteau A, Cruveiller S, Engelen S, Lajus A, Le Fèvre F, Longin C, Mornico D, Roche D, Rouy Z, Salvignol G, Scarpelli C, Thil Smith AA, Weiman M, Médigue C. 2013. MicroScope—an integrated microbial resource for the curation and comparative analysis of genomic and metabolic data. Nucleic Acids Res 41:D636–D647. <http://dx.doi.org/10.1093/nar/gks1194>.
12. Manfredi P, Pagni M, Cornelis GR. 2011. Complete genome sequence of the dog commensal and human pathogen *Capnocytophaga canimorsus* strain 5. J Bacteriol 193:5558–5559. <http://dx.doi.org/10.1128/JB.05853-11>.
13. Manfredi P, Renzi F, Mally M, Sauter L, Schmalzer M, Moes S, Cornelis GR, Jenö P, Cornelis GR. 2011. The genome and surface proteome of *Capnocytophaga canimorsus* reveal a key role of glycan foraging systems in host glycoproteins deglycosylation. Mol Microbiol 81:1050–1060. <http://dx.doi.org/10.1111/j.1365-2958.2011.07750.x>.
14. Olczak T, Sroka A, Potempa J, Olczak M. 2008. *Porphyromonas gingivalis* HmuY and HmuR: further characterization of a novel mechanism of heme utilization. Arch Microbiol 189:197–210. <http://dx.doi.org/10.1007/s00203-007-0309-7>.
15. Manfredi P, Lauber F, Renzi F, Hack K, Hess E, Cornelis GR. 2015. New iron acquisition system in *Bacteroidetes*. Infect Immun 83:300–310. <http://dx.doi.org/10.1128/IAI.02042-14>.